

Expression of Fas Ligand in Langerhans' Cell Histiocytosis: A Case Report of a Boy with Multisystem Involvement

Ching-Tien Peng,¹ Kuan-Chih Chow,^{2*} Wu-Chung Chang,³ Chang-Hai Tsai,^{1,2} Tze-Yi Lin,⁴ Sheng-Shing Lin,¹ and Chan-Fang Chiu⁵

¹Department of Pediatrics, China Medical College Hospital, Taichung, Taiwan

²Department of Medical Research, China Medical College Hospital, Taichung, Taiwan

³Department of Radiology, China Medical College Hospital, Taichung, Taiwan

⁴Department of Pathology, China Medical College Hospital, Taichung, Taiwan

⁵Department of Internal Medicine, China Medical College Hospital, Taichung, Taiwan

Previous reports of patients with Langerhans' cell histiocytosis have shown characteristics of osteolytic lesion, visceral involvement and organ dysfunction. We report a 2-year-old boy who was diagnosed as Langerhans' cell histiocytosis with a prominent hepatomegaly. X-Radiogram, computed tomography and magnetic resonance imaging revealed the osteolysis of the right iliac bone, the absence of the left eighth rib as well as the right mandible, and an enhancing mass in the left cerebellum. The data of radiography were highly suggestive of Langerhans' cell lineage. The presence of large CD1a-positive mononuclear cells associated with inflammatory cells in peripheral blood smear and bone marrow aspirate further confirmed the diagnosis. In addition, expressions of S100, CD25, CD68, CD80, CD86, and Fas ligand were identified on these cells by immunocytochemical staining. The results indicate that although these cells are activated Langerhans' cells, progression of the disease and the bone destruction could be mediated by the overt FasL expression of the cells. *Am. J. Hematol.* 61:256–261, 1999.

© 1999 Wiley-Liss, Inc.

Key words: histiocytosis; Langerhans' cell; CT; MRI; Fas ligand; immunocytochemistry

INTRODUCTION

Langerhans' cells (LCs) are major histocompatibility complex (MHC) class II-bearing bone-marrow-derived cells mainly distributed in the epidermis of the skin [1]. These cells express CD1a on the cell surface and contain racket-shaped microstructure, Birbeck granules, in the cytoplasm [2–4]. They could engulf and process antigenic material. The antigen-bearing LCs will migrate from the epidermis to the paracortex of the draining lymph node and differentiate into CD1a⁺/CD83⁺/CD86⁺ dendritic cells (DCs) or CD1a⁺/CD83⁺/CD86⁺ interdigitating reticulum cells (IDCs) [5,6]. During the differentiation, LCs could rapidly upregulate MHC class II expression and co-stimulatory factors to complete antigen presentation to T cells, whilst losing their ability to process antigens and no longer containing Birbeck granule [5,6]. The *in vitro* culture of CD34⁺ progenitor cells and LCs confirmed that CD34⁺ progenitor cells could differ-

entiate into LCs in the presence of tumor necrosis factor α (TNF α), granulocyte-macrophage colony-stimulating-factor (GM-CSF), and interleukin 4 (IL-4), and that LCs could differentiate into DCs or IDCs [7–9].

Langerhans' cell histiocytosis (LCH) is a disease characterized by the abnormal proliferation of histiocytes [10,11]. The disease can affect skin, bone, lymph node, spleen, liver, and central nervous system [12–15]. According to severity and visceral involvement of the disease, LCH has been named histiocytosis X, eosinophilic granuloma, Letterer–Siwe disease, Hand–Schüller-

Contract grant sponsor: China Medical College Hospital; Contract grant number: DMR-88-02.

*Correspondence to: Dr. Kuan-Chih Chow, Department of Medical Research, China Medical College Hospital, #2 Yuh-Der Road, Taichung, Taiwan. E-mail: t6218@hpd.cmch.org.tw

Received for publication 13 November 1998; Accepted 7 April 1999




Fig. 1. Gadolinium contrast frontal (A) and axial (B) MRIs showed osteolytic lesions of left temporal bone and left skull base together with mild enhancement of left cerebellum mass. Enhancements of the adjacent dura and sinuses suggest slow venous flow around the cerebellum mass.

Christian disease and Hashimoto–Pritzker syndrome [10–17]. All these diseases, however, share a common feature of polymorphic infiltrates consisting of mononuclear and multinuclear Langerhans' cells in the afflicted area [10–17]. Elegant study by Arenzana-Seisdedos et al. [18] showed that the purified CD1a+ cells from patients with LCH can spontaneously produce interleukin 1 (IL-1) and prostaglandin E2 (PGE2). Lader and Flanagan [19] further show that the exogenous PGE2, IL-1, and TNF α can increase human osteoclast formation and bone resorption in vitro. Although the activated LCs and osteoclast may provide partial clue to the bone lesion of LCH, serum calcium level in these patients did not change markedly. Cause of osteolytic lesion in LCH remains to be determined.

In this report, we describe a 2-year-old boy with LCH in whom the disease manifestation was noted by the radiological evaluation with marked osteolysis of the right iliac bone, the left eighth rib, and the right mandible, as well as an enhancing mass in the left cerebellum. We further used immunocytochemical method to study the expressions of CD1a, S100, lysozyme, CD14, CD25, CD68, CD80, CD86, and Fas ligand (FasL) in the peripheral blood smear as well as bone marrow aspirate. Expressions of CD1a, S100, CD25, CD68, CD80, CD86, and FasL were identified on the Langerhans' cells.

CASE REPORT

A 2-year-old boy was referred to our hospital with a history of recurrent scalp seborrheic dermatitis, bilateral otorrhea, as well as right mandibular and right hip swellings. Although no pain was complaint, a decreased will-

ing to walk was noted. Otoneurological examination, on the other hand, was within normal limits, and audiological testing on both sides was intact. Scintigrams with 99mTc indicated the increased radioactivity in right mandible, right iliac, and right acetabulum. Abdominal echonograms revealed hepatomegaly, mild dilatation of left renal pelvis, and cavity lesion of right renal parenchyma. Radiological evaluation of head and neck by magnetic resonance imaging (MRI) and computed tomography (CT) showed osteolytic lesions of left temporal bone and left skull base together with a gadolinium contrast [20] mass in the left cerebellum (Fig. 1A and B). Erosion of the right mandible (Fig. 2A) with a soft tissue mass filling (Fig. 2B) was noted by X-ray radiography and CT. A similar condition was found in the inferior portion of the right iliac bone (Fig. 3A, B, and C). Surprisingly, although X-ray and CT indicated the complete absence of the left eighth rib, no soft tissue replacement was found (Fig. 4A and B). Laboratory evaluation revealed hemoglobin of 11.9 g/dl, and white blood cell count of 34,130/ μ l with 73.4% neutrophils, 17.9% lymphocytes, 8.8% monocytes, and 0.1% eosinophils. Platelet count was 506×10^3 / μ l. Serum electrophoresis and immunoelectrophoresis were normal, as were blood chemistries (calcium 10.0 mg/dl, glucose 139 mg/dl, ALT 23 U/l, and AST 12 U/l).

METHODS

Immunocytochemical study of CD1a, S100, lysozyme, CD14, CD25, CD68, CD80, CD86 (Dako, Carpinteria, CA), and Fas ligand (FasL, Transduction Laboratories, Lexington, KY) was performed on peripheral blood




Fig. 2. (A) Frontal view of the skull by X-ray radiogram revealed osteolytic lesion of the right mandible (arrow). (B) CT image showed destruction of the right mandible and filling of the soft tissue mass (arrow).

smear and bone marrow aspirate by using the labeled Streptavidin-biotin (LSAB) method (Dako). The chromogenic reaction was visualized by peroxidase-conjugated streptavidin and aminoethyl carbazole (Sigma, St. Louis, MO). Slides were counter-stained with methyl green. Positive staining was recognized under the microscope as crimson granules. For the double staining, fluorescein-labeled antibodies (Benton-Dickinson, San Jose, CA) were added after the specimen was counter-stained. To detect viral gene product, immunocytochemistry with antibodies to LMP1, EBNA-2, and ZEBRA of Epstein-Barr virus (EBV) or to VP2 capsid protein of parvovirus B19 (PVB19) (Dako) or to late antigen of cytomegalovirus (CMV) (Novocastra, Newcastle-upon-Tyne, UK) was performed. In situ hybridization procedure [21] and polymerase chain reaction technique [22] were described previously. Probes synthesized by BRL (Bethesda, MD):

5'-fluorescein-GAGCTTATATAAGCCGAAAAAC-GTCTGAGATTCTCA-3', 5'-fluorescein-ATACTTCA-CTCCAGAAAGCAGGACAATTC-CATAGGTG-3', and 5'-fluorescein-GAGCTGAAA-CAT-CCCCAGGCCCAATCCTACCAAGGCAACC-3' were used to detect FasL expression. Chromogen was NBT/BCIP (Boehringer Mannheim, Germany). Cells from five cases of high-grade lymphoma without bone marrow involvement were used as controls.

RESULTS

Immunophenotyping on peripheral blood leukocytes was performed by using flow cytometry (FACSCalibur, Benton-Dickinson), showing the following results: Pan B (CD20) 56%; pan T (CD3) 45%, (CD4) 32%, (CD8) 13%, (CD14) 6.2%, (CD68) 7.6%, and (CD1a) 0.2%. In the peripheral blood smear, the frequency of CD68+ cells was 9.2%, and that of CD1a+ cells (Fig. 5A) was less than 1.0%. Interestingly, in addition to CD68+ cells, co-expression of FasL+ (Fig. 5B) was detected on most of the CD1a+ cells. Expression of FasL was further confirmed with ISH in these cells, and the same results were also obtained from the bone marrow aspirate as well as bone marrow biopsy (data not shown). Results of immunocytochemistry are listed in Table I with ten cases of histiocytic necrotizing lymphadenitis (HNL) and three cases of infection-associated hemaphagocytic syndrome (IAHS) [1,11-17,23] studied at the same time.

DISCUSSION

The results presented above show that the Langerhans' cells from this patient expressed CD1a and FasL. This conclusion is based on the presence of both markers on the same Langerhans' cell as detected by combining immunocytochemistry and immunofluorescence microscopy. Previous reports have described patterns of LCH with expressions of CD1a, S100, CD24, CD68, CD80, lysozyme, and Birbeck granules that suggest activated LCs in the disease [24]. Recently, by using X-chromosome inactivation assay of human androgen receptor gene, Yu et al. [25] and Willman et al. [26] indicated that LCH is a clonal histiocytic disorder. Nonetheless, they argued that although the principal feature distinguishing a neoplasm from a reactive process is monoclonality of the cell, the clonality alone was not able to fully interpret the difference in severity and progression of the disease.

From immunohistochemical data, Ruco et al. [12] suggested that the heterogeneous Langerhans' cell population in Letterer-Siwe disease could indicate the immature LCs of unknown cause. Several studies [18,27-31] supported the above hypothesis by showing that LCH correlated with functional defect of LCs in these patients.




Fig. 3. Damage of the right iliac bone. (A) LCH involving the right iliac bone was shown by X-ray; (B) CT image showed the soft tissue mass (arrow); (C) A three dimension of CT image showed the damage of the right iliac bone (arrow).




Fig. 4. Osteolytic lesion of the left eighth rib. (Row 1, A) The left eighth rib (arrow) was intact when the boy was 5 month-old. (Row 1, B) Complete destruction of the left eighth rib (arrow) was shown when the boy was 30 month-old. (Row 2, A) A three dimension CT image showed the absence of left eighth rib. (Row 2, B) Ten degree sagittal tilt of a three dimension CT image as shown in Row 2, A.

They further showed that functional defect could involve the decreased antigen-presenting activity [27,28], lack of E-cadherin expression [29], cytokine dysregulation [18,30] and the decreased number of late differentiating suppressor lymphocytes, namely CD8⁺ cells [31].

As mentioned previously, purified CD1a⁺ cells from

patients with LCH could spontaneously produce IL-1 and PGE2. Because the exogenous IL-1 and PGE2 could enhance human osteoclast function and bone resorption, there is a good reason to believe that activated LCs could induce the osteolytic lesion in LCH. Nevertheless, no hypercalcemia was detected in patients with LCH. In




Fig. 5. Expressions of CD1a and Fas ligand in the Langerhans' cell of peripheral blood smear from patient with LCH. By immunofluorescence microscopy, CD1a was detected on Langerhans' cell (A), and by immunocytochemistry, co-expression of FasL (B) was detected.

TABLE I. Immunostain Results of Patients with LCH, HNL and IAHS

Marker	LCH (n = 1)	HNL (n = 10)	IAHS (n = 3)
CD1a	+	+ ^a	+
S100	+	—	+
Lysozyme	—	+	+
CD14	+	+	+
CD25	+	+	+
CD68	+	+	+
CD80	+	+	—
CD86	+	+	—
Fas ligand	+	+	+ ^b —
CMV	—	—	+ ^b /—
EBV	—	+	—
PVB19	—	—	+

^aCD1a⁺ signals were identified on sinusoid cells.

^bSignals of FasL and CMV were only detected in a case with AIDS.

addition, our results showing the absence of left eighth rib without LC mass filling and bone pain in the patient would argue that increased osteoclast activity is solely responsible for the bone damage. Recent studies by Kawakami et al. [32] and Jilka et al. [33] clearly demonstrate that human osteoblast expresses Fas (Apo-1/CD95), and interaction between Fas and FasL could mediate osteoblast apoptosis. Constitutive expression of FasL was also detected on multiple myeloma cells that were frequently associated with osteolytic lesion and hypercalcemia [34,35]. It is worth noting that from the "local coupling" theory of bone formation and resorption the intact osseous configuration is kept by the concerted activities of osteoblast and osteoclast [32,33]. Although a malfunction of either cell type may jeopardize rigidity and structure of the bone, the complete damage requires the increased activity of osteoclast and the decreased function of osteoblast. The study of this case considered together the others [2–19] clearly indicates that LCs could express IL-1, PGE2 and FasL at the same time. The

exact role of each feature in producing the syndrome, however, remains to be determined.

ACKNOWLEDGMENTS

We thank Mr. Wen-Ji Lo for detecting CD1a⁺ cells in the peripheral blood by flow cytometry. The authors are indebted to the Section of Transfusion Medicine, Department of Laboratory Medicine and Pathology, China Medical College Hospital Taichung, Taiwan for identifying cases for this study.

REFERENCES

1. Luken JN. Langerhans' cell histiocytosis. In Lee GR, Bithell TC, Foerster J, Athens JW, Lukens JN, eds. *Wintrobe's Clinical Hematology*, Volume 2. Philadelphia: Lea & Febiger, 1993.
2. Beckstead JH, Wood GS, Turner RR. Histiocytosis X cells and Langerhans cells: enzyme histochemical and immunologic similarities. *Hum Pathol* 1984;15:826.
3. Fithian E, Kung P, Goldstein G, Rubinfeld M, Fenoglio C, Edelson R. Reactivity of Langerhans cells with hybridoma antibody. *Proc Natl Acad Sci USA* 1981;78:2541.
4. Robb IA, Jimenez CL, Carpenter BF. Birbeck granules or Birbeck junctions? Intercellular "zipperlike" lattice junctions in eosinophilic granuloma of bone. *Ultrastruct Pathol* 1992;16:423.
5. Takahashi K, Asagoe K, Zaishun J, et al. Heterogeneity of dendritic cells in human superficial lymph node: in vitro maturation of immature dendritic cells into mature or activated interdigitating reticulum cells. *Am J Pathol* 1998;153:745.
6. Strobl H, Scheinecker C, Riedl E, et al. Identification of CD68+lin-peripheral blood cells with dendritic precursor characteristics. *J Immunol* 1998;161:740.
7. Luft T, Pang KC, Thomas E, et al. A serum-free culture model for studying the differentiation of human dendritic cells from adult CD34⁺ progenitor cells. *Exp Hematol* 1998;26:489.
8. Herbst B, Kohler G, Mackensen A, Veelken H, Lindemann A. GM-CSF promotes differentiation of a precursor cell of monocytes and Langerhans-type dendritic cells from CD34⁺ haemopoietic progenitor cells. *Br J Haematol* 1998;101:231.
9. Geissmann F, Prost C, Monnet JP, Dy M, Brousse N, Hermine O. Transforming growth factor beta 1, in the presence of granulocyte/macrophage colony-stimulating factor and interleukin 4, induces differentiation of human peripheral blood monocytes into dendritic Langerhans cells. *J Exp Med* 1998;187:961.
10. Malone M. The histiocytoses of childhood. *Histopathology* 1991;19:105.
11. Cline MJ. Histiocytes and histiocytosis. *Blood* 1994;84:2840.
12. Ruco LP, Remotti D, Monardo F, et al. Letterer-Siwe disease: immunohistochemical evidence for a proliferative disorder involving immature cells of Langerhans lineage. *Virchows Arch A Pathol Anat Histopathol* 1988;413:239.
13. Nagaoka S, Maruyama R, Koike M, et al. Cytology of Langerhans' cell histiocytosis in effusions. *Acta Cytol* 1996;40:563.
14. Wang WS, Liu JH, Yang CF, Yang AH, Chen PM. Langerhans' cell histiocytosis (histiocytosis-X) in a woman with typical ultrastructure of Birbeck's granule: a case report. *Chin Med J (Taipei)* 1996;58:281.
15. Angeli SI, Luxford WM, Lo WWM. Magnetic resonance imaging in the evaluation of Langerhans' cell histiocytosis of the temporal bone: case report. *Otolaryngol Head Neck Surg* 1996;114:120.
16. Bergmann M, Yuan Y, Palm KV, Rohkamm R. Solitary Langerhans' cell histiocytosis lesion of the parieto-occipital lobe: a case report and review of the literature. *Clin Neurol Neurosurg* 1997;99:50.

17. Favara BE. Langerhans' cell histiocytosis pathobiology and pathogenesis. *Semin Oncol* 1991;18:3.
18. Arenzana-Seisdedos F, Barbey S, Virelizier JL, Kornprobst M, Nezelof C. Histiocytosis X. Purified (T6+) cells from bone granuloma produce interleukin 1 and prostaglandin E2 in culture. *J Clin Invest* 1986;77:326.
19. Lader CS, Flanagan AM. Prostaglandin E2, interleukin 1 alpha, and tumor necrosis factor-alpha increase human osteoclast formation and bone resorption in vitro. *Endocrinology* 1998;139:3157.
20. Rosenfield NS, Abrahams J, Komp D, Brain MR in patients with Langerhans cell histiocytosis: findings and enhancement with Gd-DTPA. *Pediatr Radiol* 1990;20:433.
21. Chang YT, Liu HN, Chow KC, Chen KY, Wong CK. Detection of Epstein-Barr virus in primary cutaneous amyloidosis. *Br J Dermatol* 1997;136:823.
22. Chow KC, Nacilla JQ, Witzig TE, Li CY. Is persistent polyclonal B lymphocytosis caused by Epstein-Barr virus? - A study with polymerase chain reaction and in situ hybridization. *Am J Hematol* 1992;41:270.
23. Kikuchi M. Lymphadenitis showing focal reticulum cell hyperplasia with nuclear debris and phagocytes. *Acta Hematol Jpn* 1972;35:379-380.
24. Emile J, Fraitag S, Leborgne M, Prost YD, Brousse N. Langerhans' cell histiocytosis cells are activated Langerhans' cells. *J Pathol* 1994;174:71.
25. Yu RC, Chu C, Buluwela L, Chu AC. Clonal proliferation of Langerhans cells in Langerhans cell histiocytosis. *Lancet* 1994;343:767.
26. Willman CL, Busque L, Griffith BB, et al. Langerhans'-cell histiocytosis (histiocytosis X)—a clonal proliferative disease. *N Engl J Med* 1994;331:154.
27. Yu RC, Alaibac M, Chu AC. Functional defect in cells involved in Langerhans cell histiocytosis. *Arch Dermatol Res* 1995;287:627.
28. Hosmalin A, McIlroy D, Autran B, et al. Imbalanced "memory" T lymphocyte subsets and analysis of dendritic cell precursors in the peripheral blood of adult patients with Langerhans cell histiocytosis. *Clin Exp Rheumatol* 1997;15:649.
29. Geissmann F, Emile JF, Andry P, et al. Lack of expression of E-cadherin is associated with dissemination of Langerhans' cell histiocytosis and poor outcome. *J Pathol* 1997;181:301.
30. Schultz C, Klouche M, Friedrichsdorf S, Richter N, Kroehnert B, Bucskey P. Langerhans cell histiocytosis in children: does soluble interleukin-2-receptor correlate with both disease extent and activity? *Med Pediatr Oncol* 1998;31:61.
31. Bove KE, Hurtubise P, Wong KY. Thymus in untreated systemic histiocytosis X. *Pediatr Pathol* 1985;4:99.
32. Kawakami A, Eguchi K, Matsuoka N, et al. Fas and Fas ligand interaction is necessary for human osteoblast apoptosis. *J Bone Miner Res* 1997;12:1637.
33. Jilka RL, Weinstein RS, Bellido T, Parfitt AM, Manolagas SC. Osteoblast programmed cell death (apoptosis): modulation by growth factors and cytokines. *J Bone Miner Res* 1998;13:793.
34. Villunger A, Egle A, Marschitz I, et al. Constitutive expression of Fas (Apo-1/CD95) ligand on multiple myeloma cells: a potential mechanism of tumor-induced suppression of immune surveillance. *Blood* 1997;90:12.
35. Bataille R, Chappard D, Klein B. The critical role of interleukin-6, interleukin-1B and macrophage colony-stimulating factor in the pathogenesis of bone lesions in multiple myeloma. *Int J Clin Lab Res* 1992;21:283.